


DATI PROGRAM JE ORJENTACIONI. **PRATITI IZVEŠTAVANJA PIS** (PROGNOZNO IZVEŠTAJNE SLUŽBE)!


BOLESTI I ŠTETOČINE	MART	APRIL	MAJ	JUN	JUL	AVGUST	SEPTEMBAR
BOLESTI IZDANAKA							
BOLESTI LISTA							
BOLESTI PLODA							
LISNE VAŠI, GRINJE							
MALININA BUBA, CVETOJEDI							
MUŠICA							
DROZOPHILA SUZUKII							
MUVE GALICE							



1) Neposredno pred kretanje vegetacije

Plavi preparati
Mineralno ulje


00-03



4) Pojava cvetnih pupoljaka

Kingbo AS 0,2%
Chitosan 0,2%


51-59



7) Rast plodova


Chitosan 0,2%

71-79



10) Nakon berbe


Vegard AS 0,3-0,4%



2) Otvaranje pupoljaka početak listanja

Kingbo AS 0,2%
Vegard AS 0,2-0,4%


03-09



5) Početak cvetanja

Kingbo AS 0,2%
Aposthole 0,2%


60-61



8) Sazrevanje plodova

Kingbo AS 0,2%
Perperin 0,2%
Chitosan 0,2%

81



3) Intenzivni rast listova

Aposthole 0,2%

10-31

6) Precvetavanje kraj cvetanja

Vegard AS 0,2-0,4%

67-69

9) Berba zreli plodovi (po potrebi)

Kingbo AS 0,2%
Perperin 0,2%
Chitosan 0,2%

87-89

SREDSTVA U PROGRAMU IMAJU PREVENTIVNO DEJSTVO, TREBA IH KORISTITI PRI POJAVI PRVIH SIMPTOMA. RADI BOLJE EFIKASNOSTI, OBAVEZNO KORISTITI ATOMIZERE I TRETMAN ZAPOČETI ODOZDO (NALIČJE)

* KOD POJAVE ERIOFIDNIH GRINJA KINGBO PO POTREBI PONOVI NAKON 7 DANA

NAPOMENA: **OBAVEZNO PREVENTIVNO TRETIRANJE U FAZI ZRENJA NA SVAKIH 5-7 DANA KINGBO-m (DROZOPHILA SUZUKII).**

NAKON BERBE VEGARD 0,3-0,4% (DIDYMELLA APPLANATA).