



Asian Journal of
Plant Pathology

ISSN 1819-1541



Academic
Journals Inc.

www.academicjournals.com



Review Article

Chitosan for Eco-friendly Control of Plant Disease

Oliul Hassan and Taehyun Chang

Division of Ecology and Environmental System, College of Ecology and Environmental Sciences, Kyungpook National University, 37224 Sangju-si, Gyeongsangbuk-do, Republic of Korea

Abstract

Due to the high environmental risk of chemical pesticides, biological control of plant diseases with bio-pesticides are highly encouraged and recommend. Chitosan and oligochitosan are the well-known biological control agent for its nontoxic, biodegradable and biocompatible properties. Chitosan is considered as the most abundant natural polymer with a dual effect: It controls pathogenic microorganisms by preventing growth, sporulation, spore viability, germination and disrupting cell and inducement of different defense responses in host plant inducing and/or inhibiting different biochemical activities during the plant-pathogen interaction. Chitosan has been assayed for control numerous pre and post-harvest diseases of many crops. Chitosan also has the positive effect of enriching biodiversity in the rhizosphere. Meteorological effect on chitosan is little evaluated. For achieving the goal of sustainable agriculture, chitosan will become a popular plant protectant.

Key words: Chitosan, ecofriendly, biopesticides, resistance, biodiversity

Received: December 19, 2016

Accepted: February 24, 2017

Published: March 15, 2017

Citation: Oliul Hassan and Taehyun Chang, 2017. Chitosan for eco-friendly control of plant disease. *Asian J. Plant Pathol.*, 11: 53-70.

Corresponding Author: Taehyun Chang, Division of Ecology and Environmental System, Kyungpook National University, 37224 Sangju-si, Gyeongsangbuk-do, Republic of Korea Tel: +82-54-530-1204 Fax: +82-54-530-1209

Copyright: © 2017 Oliul Hassan and Taehyun Chang. This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Chemical pesticide is of increasing concern as their negative impact on human health and environment^{1,2}. The broad spectrum pesticides may cause problems by targeting beneficial organisms and repeated use of these synthetic pesticides leads to loss of biodiversity^{3,4}. Some pesticides are not biodegradable and persistence in environment from few to many years for these reasons, they are responsible for soil, water and a wide range of environmental pollution⁵⁻⁷. Some pesticides bio-accumulated in the food chains which is in consequence affecting the human health^{8,9}. There is lots of evidence of the emerging resistance to fungicides due to the indiscriminate use of synthetic fungicides. In these circumstances control of pest with the bio-pesticides is the better alternative to the synthetic ones as the bio-pesticides have good efficacy, low residues and little or no toxicity to non-target organisms and human, causes low environmental pollution and several other advantages¹⁰.

Chitin and its derivative, chitosan have been known worldwide for its potential as a biocontrol agent. After cellulose, the second most abundant and important natural biopolymer is chitin in the world¹¹. Chitin has a variety of applications such as in medical science, chemical science and agricultural industry. In medical science, chitin has been used as a wound dressing material, drug carrier and anti-cancer drug etc., while in chemical science it has been used to prepare an affinity chromatography column and to immobilized enzyme etc.¹¹. It has been used as a food additive and bio-pesticides in the agricultural industries¹². The most important properties of chitosan such as biodegradability, nontoxicity, biocompatibility and fungicidal effect^{11,13}.

A lot of biological control agents, including natural product (derived from microorganism, higher plant and animal) and microorganisms (fungi, bacteria, virus nematodes and protozoa)^{10,14} have been used to control the plant diseases and to inhibit the growth of plant pathogens. Chitosan has some advantages over other biocontrol agent not only its, potential to control plant diseases but also its ability to induce resistance in the host plants¹⁵ and enhance biodiversity in the rhizosphere¹⁶.

The objectives this study were to summarized the efficacy of chitosan and oligo-chitosan on controlling plant diseases, mechanism of induction resistance in plant and influence on microbial community in rhizosphere.




Fig. 1(a-b): Structures of (a) Chitin and (b) Chitosan¹²

CHEMICAL STRUCTURE AND FUNCTION OF CHITOSAN

Chitosan is the derivatives of chitin. It is prepared by deacetylation of chitin to varying degree, although the deacetylation is almost never complete. When the deacetylation of chitin is reaching below 50% (depending on the origin of chitin), the product is called chitosan, it became soluble in acidic aqueous media^{11,17}. To convert chitin to chitosan, chitin is N-deacetylated by enzymatic preparation or chemical process. In the chemical process, acids or alkalis (NaOH) are used to deacetylate chitin^{17,18}. In case of enzymatic preparation, chitin deacetylases (EC 3.5.1.41) is used to catalyze the hydrolysis of N-acetamido bonds in chitin to convert it to chitosan¹⁷. The structures of chitin and chitosan¹² are shown in Fig. 1.

Due to biological (biodegradable, nontoxic and biocompatibility) and chemical (reactive amino group, hydroxyl groups, positive charge, etc.)¹⁹ properties, the chitosan has a broad range of industrial applications.

Table 1 summarizes the main application of chitosan in different industries. In this text, we will give emphasis on the use of chitosan in agriculture as elicitors, bio fungicides and modifier of plant-microbial interaction etc.

In vitro ANTIMICROBIAL ACTIVITY

***In vitro* antifungal activity:** There are several techniques that have been used to understand the antimicrobial activities of chitosan. *In vitro* antifungal activities of chitosan against several genera of fungi have been investigated and documented by several studies. Most of the study revealed that the chitosan affects mycelium growth, sporulation, morphology and molecular organization of fungus.




Fig.2(a-b): Effect chitosan preparation concentrations on the rate of mycelial growth and inhibition in V-8 juice agar medium amended with SH-1 (75 µg a.i. mL⁻¹ total glucosamine to 1,500 µg a.i. mL⁻¹ in V-8 juice agar) and SH-2 (37.5 µg a.i. mL⁻¹ total glucosamine to 750 µg a.i. mL⁻¹ in V-8 juice agar) after 10 days of inoculation, (a) Effect of mycelial radial growth and (b) Control value of mycelial radial growth⁴³

Table 1: Potential application of chitosan in different field

Major field	Potential applications
Medical and pharmaceutical	Hydrophobic drug carrier, encapsulating nutrients, fibers, fabrics, artificial organ, membrane, burn treatment, wound dressing, corneal contact lens, etc. ²⁰⁻²²
Agriculture	Laminated film, seed coating, potting mixture, fungicides, elicitors, etc. ^{23,24}
Water and waste treatment	Heavy metal trapper, odor reducer, flocculant, etc. ^{25,26}
Biotechnology	Emulsifier, chelator, immobilize and pattern biomolecules, graphitic carbon nanocapsules, tungsten carbide, etc. ²⁷
Food and beverages	Food additive, preservative, protective coating for fruits, food dyes and nutrients, etc. ²⁸⁻³⁰
Cosmetics and toiletries	Lotions, cream, nail lacquers, hair additives soap, cleansing milk, face peel, lipstick, foot powder, tooth pest and mouth wash ³¹

Chitosan effectively inhibits radial mycelium growth, sporulation, spore germination and elongation of *Fusarium* spp.,³²⁻³⁴ *Rhizopus* spp.,^{35,36} *Phytophthora* spp.,^{37,38} *Botrytis* spp.,^{39,40} and *Alternaria* spp.,⁴¹ etc. Meng *et al.*⁴² demonstrated the inhibition rate of mycelium growth of fungi depend on the concentration of chitosan and its derivatives, oligochitosan. Growth inhibition of mycelium increases with the increasing concentration of chitosan preparation SH-1 (75 µg a.i. mL⁻¹ total glucosamine to 1,500 µg a.i. mL⁻¹ in V-8 juice agar) and SH-2 (37.5 µg a.i. mL⁻¹ total glucosamine to 750 µg a.i. mL⁻¹ in V-8 juice agar) after 10 days of inoculation (Fig. 2)⁴³.

In almost all cases, spore-producing fungi produce lower spore when treated with chitosan than untreated fungi. Moreover, in some reports complete sporulation inhibition was observed after treating with chitosan. In contrary to sporulation inhibition, chitosan stimulates sporulation of *Penicillium digitatum* and *Alternaria alternata*^{44,45}. The researchers explained this phenomenon by the responses of these fungi against the chitosan induced stress.

The inhibitory effect of chitosan depends on the type of solvent. Chitosan dissolved in lactic acid show the best inhibitory effect as compared to dissolved in formic acid and acetic acid³³. Some studies have reported the antifungal

activities of chitosan depend on the incubation period. The growth inhibition of *F. oxysporum* f. sp. *radicis-lycopersici* decreases with the increased incubation period when treated with chitosan at low concentration (1.0 and 2.0 mg mL⁻¹)⁴⁶ while for *Aspergillus niger*, incubation time is very important for chitosan efficacy⁴⁷. Efficacy of chitosan also depended on the type of chitosan. Low Molecular Weight (LMW) chitosan has the highest inhibitory effect on *Rhizopus stolonifera*, *Botrytis cinerea* and *Penicillium expansum*^{39,48} while High Molecular Weight (HMW) chitosan show better efficacy on *F. oxysporum* f. sp. *vasinfectum*, *Alternaria solani* and *Valsa mali*⁴⁹. All fungus species are not similarly sensitive to chitosan³⁵. Chitosan (0.5 and 1.5%) appears to be more effective in inhibiting mycelium growth of *F. oxysporum* than *P. digitatum* and *R. stolonifera*⁴⁴. The chitosan could be formulated and applied as a natural antifungal agent into nanoparticles form, to enhance its antifungal activity. Ing *et al.*⁵⁰ reported that chitosan nanoparticles prepared from low molecular weight and high molecular weight chitosan have better inhibitory activity against *Candida albicans*, *F. solani*. Chitosan silver nanoparticles inhibit conidia germination of *Colletotrichum gloeosporioides* more effectively than chitosan alone⁵¹. Special characters of the chitosan nanoparticles such as small and compact particle




Fig. 3(a-f): Effect of chitosan on fungal morphology (a) Hyphal agglomeration, (b-c) Abnormal shapes, (d-e) Large vesicles in the mycelia (arrows) and (f) Normal hyphal growth in control³⁴

as well as high surface charge are responsible for higher antifungal activities than sole chitosan polymer⁵⁰.

Effect of chitin, chitosan and its oligomers on fungal morphology is well documented. Scanning electron microscope analysis reveal that inhibition of fungal growth in response to chitosan and its oligomers was accompanied with excessive mycelial branching, abnormal shapes, swelling and hyphae size reduction in *Fusarium* spp.^{32,34,46,52}, *B. cinerea*, *R. stolonifera*, *A. alternate* and *Penicillium expansum*^{35,53,54}. Chitosan also responsible for cytological alteration, protoplasm dissolution and large vesicle of fungus³⁴ shown in Fig. 3.

***In vitro* antibacterial activity:** As like as fungi, bacteria also highly sensitive to the chitosan and its derivatives. Most of the chitosan antibacterial reports are on human disease causing bacteria such as *Escherichia coli*, *Staphylococcus aureus* and certain *Bacillus* species. Since chitosan shows the bactericidal activity against a variety of human disease-causing bacteria, it could be expected that chitosan

may protect the plant from bacterial diseases. Recently, some researchers revealed that chitosan has strong *in vitro* antibacterial activities against plant pathogenic bacteria such as *Streptomyces scabies*⁵⁵, *Xanthomonas*⁵⁶, *Pseudomonas syringae*⁵⁷, *Pseudomonas fluorescens*⁵⁸, *Agrobacterium tumefaciens*⁴¹, *Erwinia*⁴¹, *Burkholderia seminalis*⁵⁹, *Acidovorax citrulli*⁶⁰ and *Acidovorax avenae*⁶¹. Inhibitory activity of chitosan against bacteria depends on, concentration^{41,61} molecular weight of chitosan⁶² type of bacteria (Gram-positive and Gram-negative)⁶³, type of solvent⁴¹, incubation period and abiotic factors⁶⁴. From these studies, it is clear that chitosan could be used as potential bacteria controlling agent.

***In vitro* antiviral activity:** Despite chitosan is not being a component of the virus, chitosan has been shown to control plant viral diseases. Suppression of viral infection was reported with the tobacco mosaic and necrosis viruses, potato virus X, peanut stunt virus, cucumber mosaic virus and alfalfa mosaic virus⁶⁵⁻⁶⁷. It is hypothesized that chitosan can inhibit

viral infection by inducing hypersensitive response, decreasing viability of phage bacterial cells, neutralizing the infectivity of phage particle and blocking the virulent phage replication⁶⁸.

In vitro nematocidal and anti-protozoal activity:

Nematicidal as well as the anti-protozoal activity of chitosan and its derivatives has not been sufficiently studied. From the 1980s onward, a number of studies suggested that chitosan is effective to control plant pathogenic nematode population⁶⁹⁻⁷¹. Chitinous amendments result in significant reduction in egg hatching, the viability of larvae and adult of the plant pathogenic nematode species *Meloidogyne arenaria*⁶⁸, *Meloidogyne incognita*⁷², *Meloidogyne javanica*⁷³ and *Heterodera glycines*⁷⁰. The nematocidal activity of chitosan depends on the concentration, molecular weight of chitosan. Khalil and Badawy⁷² suggested that the chitosan at low molecular weight may serve as a natural nematicide. A few studies have demonstrated that application of chitosan biomolecule is promising option for treatment trichomoniasis^{74,75}. Deacetylated chitosan oligosaccharide significantly decreases the viability of *Trichomonas vaginalis*⁷⁴ and *Trichomonas gallinae*⁷⁵. Still, a long way to go to make conclusive statements on chitosan as the antiprotozoal agent.

USE OF CHITOSAN FOR CONTROLLING PLANT DISEASE (In vivo STUDY)

Control of *Fusarium* spp.: *Fusarium* spp., is a very important plant pathogen causes disease in a wide range of plant, leading significant economic loss and mycotoxin in food and animal feed. At least 81 of 101 economically important crops are affected by *Fusarium* spp., infection. Chitosan provides a satisfactory level of plant-protection against *Fusarium* spp., as compared to chemical pesticide^{33,76,77}. To control pathogens, chitosan has been utilized in seed, soil amendment and as foliar treatment⁷⁸. *Fusarium oxysporum* the causal agent of crown and root rot disease of tomato. Foliar spraying of chitosan at concentrations ranging from 0.5-2 mg mL⁻¹ markedly reduces the number of root lesions caused by the *F. oxysporum*⁷⁹. Bhaskara Reddy *et al.*⁷⁸ reported that chitosan treatment (2-8 mg mL⁻¹) of wheat seeds reduces seed born *F. graminearum* >50%. They also suggested that chitosan treatment increase resistance in seedling and grain yield. Soil amendment with optimal concentrations of chitosan able reduced plant wilting caused by *Fusarium* spp., especially *F. acuminatum*⁸⁰. Hadwiger and Beckman⁸¹ found that the tissue is protected from *F. solani* f. sp., *pisi*, infection when chitosan applied to pea pod tissue. *Fusarium moniliforme* is an

opportunistic microorganism associated with maize, which is able to colonize in the roots and uptake nutrient. Use of chitosan as maize seed coating agent inhibit the *F. moniliforme* growth⁸². Dry rot is an important disease of stored potato caused by *F. sulphureum*. Li *et al.*³³ reported that chitosan application at the rate of 0.5 or 1% effectively controlled the dry rot of potato tuber.

Control of *Phytophthora* spp.: According to the results of several studies, *Phytophthora* spp., is effectively controlled by chitosan. Spraying and soil drench with chitosan reduced *Phytophthora infestans*^{43,83,84}, *Phytophthora cactorum*⁸⁵ and *Phytophthora capsici*⁸⁵ diseases. Plant sprayed with N, O-carboxymethyl chitosan was shown to delay late blight development (*P. infestans*)⁸³. Cho *et al.*⁸⁴ reported that oligochitosan suppressed the development of late blight on potato with the control value of 72%. Chang and Kim⁴³ after evaluating chitosan preparations (SH-1 and SH-2) on late blight suggested that SH-1 and SH-2 can be used as the natural fungicide in organic farming to increases tuber yield and to control the late blight disease. *Phytophthora cactorum* is the causal organism for crown rot in strawberry. Effect of chitosan on crown rot disease depends on the concentration of chitosan and time between application and inoculation. Disease severity decrease as the time between treatment and inoculation increased. It may take certain time interval for the plant to install its defenses. Low concentrations of chitosan were more effective against crown rot than high concentrations. *Phytophthora* root and crown rot of bell pepper caused by *Phytophthora capsicum* were better controlled when the soil of nursery bed is amended with the chitosan (0.2% w/v)⁸⁶. *Phytophthora parasitica* is a root pathogen with a wide range of host. For example, black shank (*P. parasitica*) a devastating disease of all types of cultivated tobacco. The result of Falcon *et al.*⁸⁷ showed that partially acetylated chitosan was very effective in protecting tobacco against *P. parasitica*.

Control of *Rhizoctonia* spp. and *Sclerotinia* spp.: The potential of chitosan to suppress the diseases caused by *Rhizoctonia* spp. and *Sclerotinia* spp., has been confirmed various investigations. *Rhizoctonia solani* is a soil born plant pathogen which is responsible for damping-off, sheath rot and root rot of many plant species. Acid soluble chitosan protects rice seedling from sheath blight (*R. solani*) and causes a 66-91% inhibition lesion length⁸⁸. In other studies carried out with lettuce treated with biochikol 020 PC, (2% of chitosan) showing symptoms of damping-off (*R. solani*) was significantly lower⁸⁹.




Fig. 4: Protective activity of chitosan preparations on tomato leaf mold¹⁰³

Sclerotinia sclerotiorum is another soil born plant pathogen causing root crown and stem rot in various plant species. Application of chitosan hydrolysate at the rate of 0.2% (w/v) to carrot 3 days before inoculation reduced the size and frequency of *Sclerotinia* rot as well as induced resistance⁹⁰. Yin *et al.*⁹¹ found the similar result when oligochitosan when applied to oilseed rape 3 days before inoculation.

Control of *Pythium* spp., *Penicillium* spp. and *Puccinia* spp.:

Pythium species cause diseases in a wide range of crops. As for the example root rot of cucumber and rhizome rot disease of turmeric are the very destructive disease caused by *Pythium aphanidermatum*. El Ghaouth *et al.*⁹² investigated the biological activity of chitosan on *Pythium* rot of cucumber. It is found that chitosan (100-400 mg mL⁻¹) not only control the root rot by adversely affecting the growth of *P. aphanidermatum* but also stimulated physical barrier and enzymatic defense mechanism without affecting normal growth of cucumber (Fig. 4). Symptom of phytotoxicity also was not found even in higher concentration of chitosan. Anusuya and Sathiyabama⁹³ reported that turmeric plant treated with chitosan (0.1% w/v spraying on leaves) showed increased resistance towards rhizome rot disease caused by *P. aphanidermatum*.

Blue mould and green mould diseases (caused by several species of *Penicillium*) cause considerable economic losses on fruit and vegetable during the storing period. Coating fruits and vegetable with chitosan protect them from pathogen and increase their self-life^{39,94,95}. Tomato fruits treated with 0.5 and 1% chitosan could significantly decrease blue mould caused by *P. expansum* when stored at 2°C and the effects of chitosan on this disease were increase with the increasing concentration of chitosan³⁹. Wang *et al.*⁹⁵ found that chitosan significantly inhibits blue mold caused by *P. expansum* and natural decay on jujube fruit in concentration dependent manner. Coating citrus (*Murcott tango*) fruit with chitosan

significantly reduced postharvest decay caused by *Penicillium digitatum* and *Penicillium italicum* in respect of molecular weight (high or low)⁹⁵.

Peanut (*Arachis hypogaea*) is one of the popular and high demanding crops all over the world for its nutritional value. The study conducted on peanut infection with *Puccinia arachidis* indicated that chitosan (1000 ppm) has a good controlling effect on leaf rust disease. Chitosan reduced the number of lesions and lesion diameter when applied before inoculation⁹⁶.

Control of *Botrytis* spp., *Colletotrichum* spp. and other fungal species:

Botrytis cinerea Pers., a necrotroph plant pathogen, causes gray mould disease in over 200 plant species. Many studies showed that chitosan and its derivatives have the ability to inhibit the development of gray mould in different plant species such as cucumber⁹⁷, grapes⁹⁸ and tomato⁹⁹ etc. The chitosan has the ability decrease in the gray mould disease index from 3.5 in the control plants to 0.45 when spraying on cucumber plants before 24 h of inoculation⁹⁷. Xu *et al.*⁹⁸ evaluated the antifungal activity of grapes seed extract and chitosan, alone or combined, against fungal rot (*B. cinerea*) of the grapefruits. It is found that GSE and chitosan have synergistic effect and treatment with 1% chitosan plus 0.1% GSE also significantly reduced gray mold incidence followed by treatment with 1% chitosan coating. As a biocontrol agent chitosan also found effective against tomato gray mold. About 0.2% of chitosan solution significantly inhibited tomato gray mold⁹⁹.

Colletotrichum anthracnose (*Colletotrichum* spp.) is one of the most economically important plant disease causing serious economic loss. A wide range of crops is affected by this diseases. The studies that have been conducted to determine the effects of chitosan on *Colletotrichum* anthracnose revealed that it (chitosan) has satisfiable controlling effect. More than 60% control of anthracnose on papaya fruit is achievable when applied before *C. gloeosporioides* inoculation¹⁰⁰. Significant reduction of anthracnose disease after coating tomato fruits and grape berry with 2.5% chitosan reported by Munoz *et al.*¹⁰¹. Chitosan also induces resistance in common bean plant hence effectively protected it (common bean) from anthracnose caused by *Colletotrichum lindemuthianum*¹⁰².

We evaluated the disease control efficacy of two chitosan preparations (SH-1 and SH-2) against tomato leaf mold caused by *Fulvia fulva* was under plastic greenhouse conditions¹⁰³. We made SH-1 preparation by mixing chitosan 2 g, chitooligosaccharide 20 g, wood vinegar 20 mL, rice vinegar 40 g and SH-2 by mixing chitosan 2 g, chitooligosaccharide 20 g, wood vinegar 40 mL and rice vinegar 20 g. In this study,

Table 2: Effect of chitosan on viral infection in different 'Plant-virus' systems¹¹⁴

Plant species	Virus	Effect of chitosan	Rate of inhibition ^a
<i>Phaseolus vulgaris</i>	ALMV-L	Reduction in the number of local lesion	++++
<i>Phaseolus vulgaris</i>	TNV		++++
<i>Chenopodium quinoa</i>	TNV		++
<i>Chenopodium quinoa</i>	CMV		+++
<i>Nicotiana tabacum</i> var. Samsun NN	TMV		+++
<i>Nicotiana tabacum</i> var. Xanthi nc	TMV		++
<i>Nicotiana glutinosa</i>	TMV		++
<i>Nicotiana paniculata</i>	PSV		++
<i>Phaseolus vulgaris</i>	ALMV-S	Reduction in the number of systemically infected plants	++++
<i>Phaseolus vulgaris</i>	PSV		++++
<i>Pisum sativum</i>	ALMV		+++
<i>Pisum sativum</i>	PSV		+++
<i>Lycopersicon esculentum</i>	PVX		+++

^aRate of inhibition: ++: 20-50%, +++: 50-75%, ++++: 75-100%

we applied chitosan preparation prior to the occurrence of leaf mold on tomato plants and disease severity was rated at different days after three applications at 7 days interval. It is found that both SH-1 and SH-2 formulations displayed potential disease control activity in every case and which was comparable to synthetic thiophanate-M (Fig. 4).

Similarly chitosan potential of controlling fungal diseases such as kernel rot (*Aspergillus flavus*) in preharvest maize¹⁰⁴, root rot (*Bipolaris sorokiniana*) of wheat⁶⁵, disease of pears cause by *A. kikuchiana* and *Physalospora piricola*⁴², post-harvest pathogenic fungi (*Rhizopus stolonife*, *Aspergillus niger*) in grapes¹⁰⁵, brown rot (*Monilinia fructicola*) in peach fruit¹⁰⁶, downy mildew (*Sclerospora graminicola*) of pearl millet¹⁰⁷ and downy mildew (*Elsinoe ampelina*) and anthracnose (*Plasmopara viticola*) in grapevines¹⁰⁸ was also reported.

Control of Bacterial, viral and nematode disease:

Experimental (*in vivo*) evidence has shown the bactericidal (kill bacteria directly) and bacteriostatic (stop bacteria from growing) ability of chitosan. Soil drench with chitosan significantly reduced *Ralstonia* wilt (*Ralstonia solanacearum*) incidence by 72% when the tomato is grown in the greenhouse. Regardless of application of chitosan, it improves the normal growth of tomato¹⁰⁹.

The different species of *Xanthomonas* are responsible for the many destructive diseases in different plant species such as leaf spot of tomato, leaf blight and leaf streak in rice, bacterial blight of cotton and citrus canker. For evaluating the antibacterial effect of chitosans with different molecular weights on *X. gardneri* (leaf spot) development and tomato plants protection. Coqueiro and Di Piero¹¹⁰ sprayed chitosan (1 mg mL⁻¹) onto plants 72, 48 or 24 h before inoculation with *X. gardneri*. It is found that chitosans (low molecular weight, medium molecular weight and commercial) controlled tomato bacterial spot irrespective of molecular weight and

reduced disease severity by 70% when applied 72 h before inoculation. Foliar spray of a commercial formulation of chitosan (Armour-Zen[®]) has the ability to reduce the incidences of *Xanthomonas vesicatoria* in greenhouse and field grown tomato plants¹¹¹. The strong antibacterial activity against leaf blight (*Xanthomonas oryzae* pv. *oryzae*) and leaf streak (*Xanthomonas oryzae* pv. *oryzicola*) of rice also reported by Li *et al.*¹¹².

Broccoli (*Brassica oleracea* L. var. *italica*) is one of the popular vegetables in the world. Growing of broccoli has been seriously hampered by the bacterial head rot (*Pseudomonas fluorescens*) disease. Regardless of application time (pre or post), six different combinations of chitosan solutions significantly reduced the disease incidence and the lesion diameter of broccoli inoculated with *P. fluorescens*⁵⁶. Elexa, the commercial formulation of chitosan effectively protect cucumber from the damage caused by bacterial angular leaf spot (*P. lachrymans*)¹¹³.

To date, a few studies have reported *in vivo* viral disease control effect of chitosan. The host-virus combination, chitosan concentration and mode of its application are the chitosan efficacy determining factors. Pospieszny *et al.*¹¹⁴ demonstrated that maximum 99.5-100% alfalfa mosaic virus (ALMV) inhibition is possible when 0.01-0.1% chitosan sprayed into the surface of the leaves before (1 day) inoculation of plants. It is also demonstrated that the varying effect of chitosan on viral infection in different 'Plant-virus' systems. The highest level of inhibition of viral infection was observed in bean leaves while TMV, TNV and CMV's local infection inhibition in tobacco and *C. quinoa* leaves was not so pronounced in these host (Table 2). Induction of antiviral defensive reactions may varies with the host plant species.

Tobacco Necrosis Virus (TNV) is one of the greatest threats for the cultivation of the tobacco. Chitosan reduces the virus-induced necrotic lesions by 32- 83% when applied on tobacco at the rate of 0.1%¹¹⁵. Chitosan has the ability to

delay the viral diseases. In comparison to control plant, chitosan retarded tomato yellow leaf curl disease symptoms about 7-14 days.

A soil amendment with chitosan was shown to reduce nematode parameters, i.e., number of galls, egg masses, females/root system and number of juveniles/250 g soil. El-Sayed *et al.*⁷³ reported that the standard concentration (1:1 dilution) of LMW and HMW chitosan reduces the number of gall by 90 and 93% respectively. Nunes da Silva *et al.*¹¹⁶ also reported that application of all molecular weight chitosan decreases the severity of the Pine Wilt Disease (PWD) caused by *Bursaphelenchus xylophilus* (pine wood nematode). They also reported that HMW chitosan is more effective than LMW chitosan.

MECHANISMS OF CHITOSAN AND ITS DERIVATIVES IN REDUCING PLANT DISEASES

There are a few concrete proofs regarding the anti-microbial mechanism of chitosan and its ability to elicit plant defense responses. The growing evidence is showing

that the chitosan and its derivatives have a double mechanism of action: They inhibit the growth of the pathogen and they modify the plant defense responses^{99,117-119}. The possible mechanisms of chitosan as a bio-pesticides is summarized in Fig. 5.

It is hypothesized that the presence of positive charge and reactive amino groups (protonated amino group on the structure of chitosan) are the base of the direct antimicrobial activities of chitosan. The electrostatic interaction between positively charged chitosan and its derivatives with negatively charged microbial cell membranes resulting in agglutination, disruption and alteration of the intracellular ultrastructure of the cell membrane and thereby causing the death organism (Fig. 6)^{33,120-122}.

Chung and Chen¹²³ reported that chitosan react on both the cell membrane and the cell wall and inactivated the pathogen via a two-step sequential mechanism: An initial separation of the cell wall from its cell membrane, followed by destruction of the cell membrane. Chitosan may interfere the function of the genetic material of the pathogen. The nucleic acid (DNA/RNA) possesses the phosphate groups in the main




Fig. 5: Modes of action of chitosan and its derivatives as antimicrobial compounds and the ability to elicit natural plant defense responses¹¹⁷




Fig. 6: Mechanism of killing of bacterial cell by chitosan¹²²

chain which are negatively charged. The chitosan could penetrate cell wall¹⁷ and bind to the negatively charged DNA and alter its confirmation in consequence block the synthesis of mRNA and essential protein of pathogen^{89,121,124}. The mode of action of chitosan may also be related to its ability to chelate some essential nutrients, metal ions and trace elements necessary for the growth of bacteria and fungi¹²⁵⁻¹²⁷.

Another possible mean of antimicrobial potential of chitosan is to deposit on the pathogen surface and the formation of a dense polymer film. This dense polymer film blocks the nutrient flow and metabolism process of microbes which are essential for their survival in nature¹¹⁷.

Plant protect themselves from pathogens by developing a stunning array of structural, chemical and protein-based defenses designed to recognize invading pathogens and arrest them before they are able to cause extensive damages¹²⁸.

Some reports indicate that chitosan and its derivatives are effective elicitor and inducer of plant systemic acquired resistance to pathogens. Chitosan and oligochitosan induce hosts to produce defense-related protein^{117,124}, enzymes^{117,129} and secondary metabolites¹¹⁷. Chitosan and its derivatives increase level and activity of glucanase in rice, tobacco, wheat, etc.^{15,130,131}. It is also reported that chitosan and its derivatives increase, level and activity of chitinase, phenylalanine ammonia-lyase, peroxidase, polyphenol oxidase, catalase and superoxide dismutase in orange, tomato, cucumber, table grapes, strawberries, sweet cherries, pears and *Zanthoxylum bungeanum* (Huajiao in Chinese)^{43,117-119}. All the mentioned protein and enzymes directly or indirectly degrade the pathogen.

The genes that functionally develop disease resistance are generally called pathogenesis-related genes. Some reports showed that chitosan employs multiple modes to increase pathogenesis-related gene function. As, for example, Hoat *et al.*¹³² reported that chitosan and chitin strongly activated the expression of the pathogenesis-related gene (PR-10) in oat leave.

It is well accepted that plant secondary metabolites such as phytoalexins, lignin, suberization and phenolic compounds play a predominant role in plant defense system. The effect of chitosan on the accumulation of defense-related secondary metabolites in the plant tissue was studied from the 1980s when Hadwiger and Beckman⁸¹ found chitosan at a concentration as low as $0.9 \mu\text{g mL}^{-1}$ can induce a phytoalexin accumulation in soybean pod in 24 h. These biopolymers (chitosan and its derivatives) also have the ability to enhance production and accumulation of phenolic compound (polyphenol), lignin, suberin and callose in different host plant^{83,78,131,133,134}. Accumulation of the polyphenol is the plant part dependent phenomenon. As for example it is found that the content of total polyphenol in the stem was significantly increased by chitosan preparation application in the field (Fig. 7)⁴³.

The resistance plants have the ability of timely recognition of the plant pathogen and the rapid and fruitful activation of the defense system. The activation of defense system manifested by the Hypersensitive Response (HR) at the site of infection and activating defense in uninfected plant part¹³⁵. The development of HR and the Systemic Acquired Resistance (SAR) govern by the combined genotypes of host and pathogen (gene for gen theory) and depend on a




Fig. 7: Accumulation of the polyphenol in the plant part after application of chitosan preparation. SH-1 and SH-2 are the chitosan preparation⁴³

complex exchange of signal occurring under given environmental condition. The signal transduction pathways composed of activation of the specific receptor on the cell membrane or located intracellular followed by transmission of the signal into the target cell mediated by one or more second messengers and create a series of physiological responses. To our knowledge, only the kinase 1 (CERK1) and the lectin are the two cell surface or membrane receptors that are likely to bind chitosan and its oligomers^{136,137}. Chitosan also influences the plant's DNA conformation. Due to the high affinity for negatively charged phosphates of the DNA backbone, chitosan may alter chromatin via competition with basic nuclear proteins for DNA attachment sites that block gene transcription in turn^{124,132}.

On the basis of the result of several studies, Reactive Oxygen Species (ROS), Ca²⁺, Nitric Oxide (NO), ethylene (ET), Jasmonic Acid (JA), Salicylic Acid (SA) and abscisic acid (ABA) all involved in chitosan-mediated signal pathway^{117,138}.

It may conclude that chitosan and its derivatives can directly destroy the pathogen and activated the plant immune system through different signaling pathways or involved in signal transduction as a regulatory molecules. Despite extensive research, the mechanism and mode of action of chitosan in reducing plant diseases have not been explicated clearly. However, the mechanisms governing the mode of action of these biopolymers yet to be studied.

ECOLOGICAL CONSIDERATION

Effect of chitosan on environment: Chitosan and its derivatives have emerged as the best ecofriendly bio-pesticides for their unique plant pathogen controlling properties, safety to human and mammals and non-toxicity to

other beneficial microbes. The US Environmental Protection Agency (EPA) concluded in 2015¹³⁹ that there are no risk to the environment because chitosan has not shown toxicity in mammals and it is natural biopolymers. The National Pesticides Information Center (NPIC) database also indicates that there has been no report on environmental degradation due to the use of chitosan.

Effect of chitosan on microbial community in soils: Soil harbours most of the microorganism. Plant interact with soil microorganism in a different way that directly or indirectly influence the growth and development of the plant. Symbiotic microorganism such as mycorrhiza and rhizobia (N-fixing bacteria) promote the growth of the plant by enhancing the plant nutritional status while soil born pathogen limit the plant growth. There are other groups of microorganism which are antagonistic to soil born pathogen hence indirectly influence the plant growth. It is believed that addition of chitosan and its oligomers improves the soil structure and alter the rhizospheric environment to shift the microbial balance in favour of beneficial organisms and to the detriment of plant pathogens^{24,140}.

There is no substantial body of evidence that chitosan supplementation increases both population and efficacy of antagonists in the rhizosphere. Mulawarman *et al.*¹⁴¹ reported that bacterial population densities of bacterial genera: *Acinetobacter* and *Pseudomonas* increased by 8.9 and 81%, respectively when soil amended with chitosan.

Pastucha¹⁴² isolated the greatest number of bacteria and fungi with antagonistic effect towards plant pathogens from rhizosphere soil of soybean after using chitosan three times. *Lysobacter enzymogenes* is an antagonist biocontrol agent for managing root and crown rot in cucumber caused by *Pythium aphanidermatum* (Edson) Fitzp. The addition of chitosan (depending on concentration) in soil increases cell number (cells per gram root) on the root as well as the efficacy of *L. enzymogenes*⁴³. *Bacillus subtilis* is another, the most widely used biopesticide in agriculture which action against powdery mildew and number of population increases with the addition of chitosan in strawberry^{144,145}. Mishra *et al.*¹⁴⁶ observed the substantially higher number of bacterial cells (*Pseudomonas* sp.) in tomato roots by scanning electron microscopy after the application of *Pseudomonas* sp., in combination with chitosan. It is also observed that chitosan enhance the efficacy of *Pseudomonas* sp., against tomato leaf curl virus (ToLCV) in tomato.

Mycorrhizal fungi, the most ubiquitous of root-associated plant symbioses, enhance the nutrient and water uptake efficiency of the plant. Our previous study¹⁶ showed that




Fig. 8(a-b): Effect of chitosan on soil microbial community. Relative abundance of soil microbial groups and physiological parameter were determined by analyzing phospholipid fatty acids (PLFA) in the soils amended with chitosan powder and solution in a cucumber greenhouse at 160 days after soil amendment, (a) Mean microbial group and (b) Physiological indicator of microbes¹⁶

vesicular-arbuscular mycorrhizal fungi (VAM) significantly increased in soils amended with chitosan powder. It is also subsisted that chitosan powder changed the soil microbial community and the effects maintained up to 160 days after soil application (Fig. 8).

In legume nitrogen fixation process, a group of free living bacteria (rhizobia) colonized in the specialized root nodules. The addition of short-chain acetylated chitin derivatives can induce nodulation in *Medicago sativa* hence, influence the N-fixing bacterial population size and activity¹⁴⁷.

Effects of chitosan in stimulating beneficial soil microbial activity can be explained by following major points: (1) Chitosan may serve as nutrient (C and N) source for the microbes^{142,145}, (2) Chitosan may induce the chitinases producing and antagonistic gene expression in biological control agents^{24,143}, (3) Chitosan may induce horizontal gene transfer (natural competence) where DNA was absorbed and recombined into the chitinolytic bacterium and able to being adapted quickly to changing conditions²⁴, (4) Chitosan may help the antagonistic biological control agents to attack

plant pathogen by disrupting the formation of biofilms produced by pathogenic microbes²⁴, (5) Chitinous nod (lipochito-oligosaccharides) and Myc (signal molecules) factors in growth-promoting microbes (mycorrhiza and *Rhizobium*) sensitizes plant roots to nodule formation and to interact with mycorrhizal fungi. Chitin-derivatives which are biochemically similar to nod and myc factors may induce nodulation and the formation of root connections with nitrogen-fixing bacteria and mycorrhizal fungi^{24,148}.

The soil microbial activities are influenced not only by chitosan but also root exudates, organic matter content soil and applied fertilizer^{149,150}. Abundant occurrence of microorganisms in the soil might have been caused by the synergistic effect of chitosan and root exudates, organic matter and fertilizer. So actual effect of chitosan and its derivatives on beneficial soil microbial activities need to be discovered.

Regulation of tolerance to abiotic stresses: Abiotic stresses such as drought, soil pollution with heavy metal, heat and high and low temperature etc., adversely affect the potential production of crops. Chitosan and its derivatives based treatments can improve plant survival and performance in stressful growing environment. Chitosan increases the chlorophyll concentration and performance of photosynthesis under drought condition^{151,134}. Plant need well developed roots system to survive under scare water condition. Zeng and Luo¹⁵¹ reported that chitosan stimulates the root system development of wheat seedling and in consequence strengthen the capability of water absorption. Excessive transpiration is not desirable in dry condition. The concentration of ABA and JA have been found to increase in response to chitosan treatment and these hormones are well known for their ability to control stomatal aperture^{23,152}. Herde *et al.*¹⁵² observed that both ABA and JA reduce the transcription rate in the tomato plant. It is also reported that chitosan induces drought resistance in rice, drought tolerance apple, coffee and paper^{23,153,154}. Bittelli *et al.*¹⁵³ found that foliar application of chitosan on pepper leaves induced partial or full closure of plant stomata, resulting in reduced water use by 26-43%.

Under the adverse environmental condition, Reactive Oxygen Species (ROS) such as peroxides, superoxide, hydroxyl radical and singlet oxygen level can increase dramatically. This may cause significant damage to the lipid cell membrane, protein and nucleic acid instead of acting as a messenger in normal cell signal transduction, cell cycling and the plant immune system. The chitosan Reactive Oxygen Species (ROS) scavenging activities were investigated by Je and Kim¹⁵⁵. It is found that chitosan showed scavenging effects

against hydroxyl and superoxide anion radical, the effects were 91.67 and 65.34%, respectively. These results suggest that the scavenging effect of chitosan depends on their degree of deacetylation and substituted group. Rajalakshmi *et al.*¹⁵⁶ also reported that chitosan and oligo chitosan possess potent antioxidant activity and hydrogen peroxide and superoxide scavenging activity.

The chelating ability of chitosan and oligo-chitosan led to the interest in their potential application to the remediation of contaminated soil and water with heavy metal, dye and accidentally application of excess fertilizer and pesticides^{25,26,157-159}. The presence of amino groups (-NH) in chitosan give the vital binding site for complexing with non-nutrient elemental ions, including a number of heavy metal cations^{157,159}. In this way, chitosan protects the plant from the effects of heavy metals, excessive fertilizers and pesticides.

CONCLUSION

In the past few decades, chitosan and its derivatives based pesticides are gaining more attention because of their non-toxic, biodegradable, biocompatible and eco-friendly nature. Chitosan and its derivatives have been repeatedly shown very impressive results as bio-pesticide against many pests of agricultural importance which is comparable to those achieved with current synthetic pesticides and fertilizers. Chitosan is considered as the most abundant natural polymer with a dual effect: It controls pathogenic microorganisms by preventing growth, sporulation, spore viability, germination and disrupting cell and inducement of different defense responses in host plant inducing and/or inhibiting different biochemical activities during the plant-pathogen interaction. The plant can also acquire enhanced tolerance to the stressful growing environment after application of chitosan and oligochitosan. Along with these, the stimulating effect of chitosan on beneficial microorganism lead this natural product to incorporate in to the Integrated Pest Management (IPM) strategies.

There are several issues associated with the commercial use of chitosan based pesticides that are still to be investigated. Like photosensitivity of chitosan derivatives (amorphous chitosan derivatives may absorb UV spectrum) has to be managed for its long lasting effectiveness. The effective concentration and application method of chitosan products is not defined yet. The biochemical mode of action of chitosan and oligochitosan for controlling plant diseases are still limited and unclear. So, much of study is still to be done in this field.

SIGNIFICANCE STATEMENTS

The second most abundant and important natural biopolymer is chitin in the world. Chitin has been used in different industries such as in medicine, chemical, cosmetics and agriculture. In the agricultural industry, these biopolymers have been using different ways including biopesticides for controlling plant diseases, bio-transformant for the production of quality food products preservative to protect foods from microbial deterioration, the chelator of waste material from food processing discards, the purifier of water and fruit juices etc. Use of chitosan as biopesticides is very effective and eco-friendly because, it is biodegradable, non-toxic and biocompatible. This study presents a comprehensive review of the antifungal action of and plant protection by chitosan.

ACKNOWLEDGMENTS

This study was supported by a grant from "Cooperative research program for FTA correspondence competitiveness improvement technology development (Project No. PJ011697032016)" in rural development administration, Korea.

REFERENCES

1. Jeyaratnam, J., 1985. Health problems of pesticide usage in the third world. *Br. J. Ind. Med.*, 42: 505-506.
2. Igbedioh, S.O., 1991. Effects of agricultural pesticides on humans, animals and higher plants in developing countries. *Arch. Environ. Health*, 46: 218-224.
3. Liebich, J., A. Schaffer and P. Burauel, 2003. Structural and functional approach to studying pesticide side-effects on specific soil functions. *Environ. Toxicol. Chem.*, 22: 784-790.
4. Yasmin, S. and D. D'Souza, 2010. Effects of pesticides on the growth and reproduction of earthworm: A review. *Applied Environ. Soil Sci.* 10.1155/2010/678360
5. Bocquene, G. and A. Franco, 2005. Pesticide contamination of the coastline of Martinique. *Mar. Pollut. Bull.*, 51: 612-619.
6. Toan, V.D., V.D. Thao, J. Walder, H.R. Schmutz and T.C. Ha, 2007. Contamination by Selected Organochlorine Pesticides (OCPs) in surface soils in Hanoi, Vietnam. *Bull. Environ. Contam. Toxicol.*, 78: 195-200.
7. Firbank, L.G., S. Petit, S. Smart, A. Blain and R.J. Fuller, 2008. Assessing the impacts of agricultural intensification on biodiversity: A British perspective. *Philos. Trans. R. Soc. London B: Biol. Sci.*, 363: 777-787.
8. Ferencz, L. and A. Balog, 2010. A pesticide survey in soil, water and foodstuffs from Central Romania. *Carpath. J. Earth Environ.*, 5: 111-118.
9. Cui, L., J. Ge, Y. Zhu, Y. Yang and J. Wang, 2015. Concentrations, bioaccumulation and human health risk assessment of organochlorine pesticides and heavy metals in edible fish from Wuhan, China. *Environ. Sci. Pollut. Res.*, 22: 15866-15879.
10. Copping, L.G. and J.J. Menn, 2000. Biopesticides: A review of their action, applications and efficacy. *Pest Manage. Sci.*, 56: 651-676.
11. Rinaudo, M., 2006. Chitin and chitosan: Properties and applications. *Prog. Polym. Sci.*, 31: 603-632.
12. Zhang, H., R. Li and W. Liu, 2011. Effects of chitin and its derivative chitosan on postharvest decay of fruits: A review. *Int. J. Mol. Sci.*, 12: 917-934.
13. Kumar, M.N.V.R., 2000. A review of chitin and chitosan applications. *React. Funct. Polym.*, 46: 1-27.
14. Yoon, M.Y., B. Cha and J.C. Kim, 2013. Recent trends in studies on botanical fungicides in agriculture. *Plant Pathol. J.*, 29: 1-9.
15. Yin, H., X. Zhao and Y. Du, 2010. Oligochitosan: A plant diseases vaccine-a review. *Carbohydr. Polym.*, 82: 1-8.
16. Park, K.C. and T.H. Chang, 2012. Effect of chitosan on microbial community in soils planted with cucumber under protected cultivation. *Korean J. Hortic. Sci. Technol.*, 30: 261-269.
17. Younes, I. and M. Rinaudo, 2015. Chitin and chitosan preparation from marine sources. Structure, properties and applications. *Mar. Drugs*, 13: 1133-1174.
18. No, H.K. and S.P. Meyers, 1995. Preparation and characterization of chitin and chitosan-a review. *J. Aquat. Food Prod. Technol.*, 4: 27-52.
19. Dutta, P.J., J. Dutta and V.S. Tripathi, 2004. Chitin and chitosan: Chemistry, properties and applications. *J. Scient. Ind. Res.*, 63: 20-31.
20. Miwa, A., A. Ishibe, M. Nakano, T. Yamahira, S. Itai, S. Jinno and H. Kawahara, 1998. Development of novel chitosan derivatives as micellar carriers of taxol. *Pharmaceut. Res.*, 15: 1844-1850.
21. Liu, W., S.J. Sun, X. Zhang and K.D. Yao, 2003. Self-aggregation behavior of alkylated chitosan and its effect on the release of a hydrophobic drug. *J. Biomater. Sci. Polym. Edn.*, 14: 851-859.
22. Dai, T., M. Tanaka, Y.Y. Huang and M.R. Hamblin, 2011. Chitosan preparations for wounds and burns: Antimicrobial and wound-healing effects. *Expert Rev. Anti-Infect. Ther.*, 9: 857-879.
23. Hoagland, P.D. and N. Parris, 1996. Chitosan/pectin laminated films. *J. Agric. Food Chem.*, 44: 1915-1919.
24. Sharp, R.G., 2013. A review of the applications of chitin and its derivatives in agriculture to modify plant-microbial interactions and improve crop yields. *Agronomy*, 3: 757-793.
25. Sobahi, T.R.A., M.Y. Abdelaal and M.S.I. Makki, 2014. Chemical modification of Chitosan for metal ion removal. *Arab. J. Chem.*, 7: 741-746.

26. Zhang, L., Y. Zeng and Z. Cheng, 2016. Removal of heavy metal ions using Chitosan and modified Chitosan: A review. *J. Mol. Liquids*, 214: 175-191.
27. Hirano, S., 1996. Chitin biotechnology applications. *Biotechnol. Annu. Rev.*, 2: 237-258.
28. Sapers, G.M., 1992. Chitosan enhances control of enzymatic browning in apple and pear juice by filtration. *J. Food Sci.*, 57: 1192-1193.
29. Roller, S. and N. Covill, 1999. The antifungal properties of chitosan in laboratory media and apple juice. *Int. J. Food Microbiol.*, 47: 67-77.
30. Shahidi, F., J.K.V. Arachchi and Y.J. Jeon, 1999. Food applications of chitin and chitosans-review. *Trends Food Sci. Technol.*, 10: 37-51.
31. Jimtaisong, A. and N. Saewan, 2014. Utilization of carboxymethyl chitosan in cosmetics. *Int. J. Cosmet. Sci.*, 36: 12-21.
32. El Hassni, M., A. El Hadrami, F. Daayf, E.A. Barka and I. El Hadrami, 2004. Chitosan, antifungal product against *Fusarium oxysporum* f. sp. *albedinis* and elicitor of defence reactions in date palm roots. *Phytopathologia Mediterranea*, 43: 195-204.
33. Li, Y.C., X.J. Sun, Y. Bi, Y.H. Ge and Y. Wang, 2009. Antifungal activity of Chitosan on *Fusarium sulphureum* in relation to dry rot of potato tuber. *Agric. Sci. China*, 8: 597-604.
34. Al-Hetar, M.Y., M.A. Zainal Abidin, M. Sariah and M.Y. Wong, 2011. Antifungal activity of chitosan against *Fusarium oxysporum* f. sp. *Cubense*. *J. Applied Polym. Sci.*, 120: 2434-2439.
35. El Ghaouth, A., J. Arul, J. Grenier and A. Asselin, 1992. Antifungal activity of chitosan on two postharvest pathogens of strawberry fruits. *Phytopathology*, 82: 398-402.
36. Hernandez-Lauzardo, A.N., M.G. Velazquez-del Valle, L. Veranza-Castelan, G.E. Melo-Giorgana and M.G. Guerra-Sanchez, 2010. Effect of chitosan on three isolates of *Rhizopus stolonifer* obtained from peach, papaya and tomato. *Fruits*, 65: 245-253.
37. Xu, J., X. Zhao, X. Han and Y. Du, 2007. Antifungal activity of oligochitosan against *Phytophthora capsici* and other plant pathogenic fungi *in vitro*. *Pestic. Biochem. Physiol.*, 87: 220-228.
38. Sunpapao, A. and C. Pornsuriya, 2014. Effects of chitosan treatments on para rubber leaf fall disease caused by *Phytophthora palmivora* Butler-a laboratory study. *Songklanakarin J. Sci. Technol.*, 36: 507-512.
39. Liu, J., S.P. Tian, X. Meng and Y. Xu, 2007. Effects of chitosan on control of postharvest diseases and physiological responses of tomato fruit. *Postharvest Biol. Technol.*, 44: 300-306.
40. Silva, Jr. S., N.P. Stamford, M.A.B. Lima, T.M.S. Arnaud, M.M. Pintado and B.F. Sarmiento, 2014. Characterization and inhibitory activity of chitosan on hyphae growth and morphology of *Botrytis cinerea* plant pathogen. *Int. J. Applied Res. Nat. Prod.*, 7: 31-38.
41. Rabea, E.I. and W. Steurbaut, 2010. Chemically modified chitosans as antimicrobial agents against some plant pathogenic bacteria and fungi. *Plant Protect. Sci.*, 46: 149-158.
42. Meng, X., L. Yang, J.F. Kennedy and S. Tian, 2010. Effects of chitosan and oligochitosan on growth of two fungal pathogens and physiological properties in pear fruit. *Carbohydr. Polym.*, 81: 70-75.
43. Chang, T. and B.S. Kim, 2012. Application of chitosan preparations for eco-friendly control of potato late blight. *Res. Plant Dis.*, 18: 338-348.
44. Bautista-Banos, S., M. Hernandez-Lopez and E. Bosquez-Molina, 2004. Growth inhibition of selected fungi by chitosan and plant extracts. *Revista Mexicana Fitopatologia*, 22: 178-186.
45. Bhaskara Reddy, M.V., J. Arul, E. Ait-Barka, P. Angers, C. Richard and F. Castaigne, 1998. Effect of chitosan on growth and toxin production by *Alternaria alternata* f. sp. *lycopersici*. *Biocontrol Sci. Technol.*, 8: 33-43.
46. Benhamou, N., 1992. Ultrastructural and cytochemical aspects of chitosan on *Fusarium oxysporum* f. sp. *radicis-lycopersici*, agent of tomato crown and root rot. *Phytopathology*, 82: 1185-1193.
47. Plascencia-Jatomea, M., G. Viniegra, R. Olayo, M.M. Castillo-Ortega and K. Shirai, 2003. Effect of chitosan and temperature on spore germination of *Aspergillus niger*. *Macromol. Biosci.*, 3: 582-586.
48. Hernandez-Lauzardo, A.N., S. Bautista-Banos, M.G. Velazquez-del Valle, M.G. Mendez-Montealvo, M.M. Sanchez-Rivera and L.A. Bello-Perez, 2008. Antifungal effects of chitosan with different molecular weights on *In vitro* development of *Rhizopus stolonifer* (Ehrenb.:Fr.) Vuill. *Carbohydr. Polym.*, 73: 541-547.
49. Guo, Z., R. Chen, R. Xing, S. Liu and H. Yu *et al.*, 2006. Novel derivatives of chitosan and their antifungal activities *In vitro*. *Carbohydr. Res.*, 341: 351-354.
50. Ing, L.Y., N.M. Zin, A. Sarwar and H. Katas, 2012. Antifungal activity of chitosan nanoparticles and correlation with their physical properties. *Int. J. Biomater.* 10.1155/2012/632698
51. Chowdappa, P., S. Gowda, C.S. Chethana and S. Madhura, 2014. Antifungal activity of chitosan-silver nanoparticle composite against *Colletotrichum gloeosporioides* associated with mango anthracnose. *Afr. J. Microbiol. Res.*, 8: 1803-1812.
52. Prapagdee, B., K. Kotchadat, A. Kumsopa and N. Visarathanonth, 2007. The role of chitosan in protection of soybean from sudden death syndrome caused by *Fusarium solanif*. sp. *glycines*. *Bioresour. Technol.*, 98: 1353-1358.
53. De Oliveira, Jr. E.N., I.S. de Melo and T.T. Franco, 2012. Changes in hyphal morphology due to chitosan treatment in some fungal species. *Braz. Arch. Biol. Technol.*, 55: 637-646.
54. Alfaro-Gutierrez, I.C., M.G. Guerra-Sanchez, A.N. Hernandez-Lauzardo and M.G. Velazquez-del Valle, 2014. Morphological and physiological changes on *Rhizopus stolonifer* by effect of chitosan, oligochitosan or essential oils. *J. Phytopathol.*, 162: 723-730.

55. Beausejour, J., N. Clermont and C. Beaulieu, 2003. Effect of *Streptomyces melanosporofaciens* strain EF-76 and of chitosan on common scab of potato. *Plant Soil*, 256: 463-468.
56. Li, B., X. Wang, R. Chen, W. Huangfu and G.L. Xie, 2008. Antibacterial activity of chitosan solution against *Xanthomonas* pathogenic bacteria isolated from *Euphorbia pulcherrima*. *Carbohydr. Polym.*, 72: 287-292.
57. Ferrante, P. and M. Scortichini, 2010. Molecular and phenotypic features of *Pseudomonas syringae* pv. *actinidiae* isolated during recent epidemics of bacterial canker on yellow kiwifruit (*Actinidia chinensis*) in central Italy. *Plant Pathol.*, 59: 954-962.
58. Li, B., B.P. Liu, T. Su, Y. Wang and G. Xie *et al.*, 2010. Effect of chitosan solution on the inhibition of *Pseudomonas fluorescens* causing bacterial head rot of broccoli. *Plant Pathol. J.*, 26: 189-193.
59. Lou, M.M., B. Zhu, I. Muhammad, B. Li and G.L. Xie *et al.*, 2011. Antibacterial activity and mechanism of action of chitosan solutions against apricot fruit rot pathogen *Burkholderia seminalis*. *Carbohydr. Res.*, 346: 1294-1301.
60. Li, B., S. Shi, C. Shan, Q. Zhou and M. Ibrahim *et al.*, 2013. Effect of chitosan solution on the inhibition of *Acidovorax citrulli* causing bacterial fruit blotch of watermelon. *J. Sci. Food Agric.*, 93: 1010-1015.
61. Yang, C., B. Li, M. Ge, K. Zhou and Y. Wang *et al.*, 2014. Inhibitory effect and mode of action of chitosan solution against rice bacterial brown stripe pathogen *Acidovorax avenae* subsp. *avenae* RS-1. *Carbohydr. Res.*, 391: 48-54.
62. Liu, N., X.G. Chen, H.J. Park, C.G. Liu, C.S. Liu, X.H. Meng and L.J. Yu, 2006. Effect of MW and concentration of chitosan on antibacterial activity of *Escherichia coli*. *Carbohydr. Polym.*, 64: 60-65.
63. Shanmugam, A., K. Kathiresan and L. Nayak, 2016. Preparation, characterization and antibacterial activity of chitosan and phosphorylated chitosan from cuttlebone of *Sepia kobeensis* (Hoyle, 1885). *Biotechnol. Rep.*, 9: 25-30.
64. Chung, Y.C., H.L. Wang, Y.M. Chen and S.L. Li, 2003. Effect of abiotic factors on the antibacterial activity of chitosan against waterborne pathogens. *Bioresour. Technol.*, 88: 179-184.
65. Badawy, M.E.I. and E.I. Rabea, 2011. A biopolymer chitosan and its derivatives as promising antimicrobial agents against plant pathogens and their applications in crop protection. *Int. J. Carbohydr. Chem.* 10.1155/2011/460381
66. Nagorskaya, V., A. Reunov, L. Lapshina, V. Davydova and I. Yermak, 2014. Effect of chitosan on Tobacco Mosaic Virus (TMV) accumulation, hydrolase activity and morphological abnormalities of the viral particles in leaves of *N. tabacum* L. cv. Samsun. *Virologica Sinica*, 29: 250-256.
67. Jia, X., Q. Meng, H. Zeng, W. Wang and H. Yin, 2016. Chitosan oligosaccharide induces resistance to *Tobacco mosaic virus* in *Arabidopsis* via the salicylic acid-mediated signalling pathway. *Scient. Rep.*, Vol. 6. 10.1038/srep26144
68. Chirkov, S.N., 2002. The antiviral activity of chitosan (review). *Applied Biochem. Microbiol.*, 38: 1-8.
69. Mian, I.H., O. Godoy, R.A. Shelby, R. Rodriguez-Kabana and G. Morgan-Jones, 1982. Chitin amendments for control of *Meloidogyne arenaria* in infested soil. *Nematropica*, 12: 71-84.
70. Rodriguez-Kabana, R., G. Morgan-Jones and B.O. Gintis, 1984. Effects of chitin amendments to soil on Heterodera glycines, microbial populations and colonization of cysts by fungi. *Nematropica*, 14: 10-25.
71. Kalaiarasan, P., P. Lakshmanan, G. Rajendran and R. Samiyappan, 2006. Chitin and chitinolytic biocontrol agents for the management of root knot nematode, *Meloidogyne arenaria* in groundnut (*Arachis hypogaea* L.) cv. Co3. *Indian J. Nematol.*, 36: 181-186.
72. Khalil, M.S. and M.E.I. Badawy, 2012. Nematicidal activity of a biopolymer chitosan at different molecular weights against root-knot nematode, *Meloidogyne incognita*. *Plant Protect. Sci.*, 48: 170-178.
73. El-Sayed, S.M. and M.E. Mahdy, 2015. Effect of chitosan on root-knot nematode, *Meloidogyne javanica* on tomato plants. *Int. J. ChemTech Res.*, 7: 1985-1992.
74. Shin, W.S., J.C. Kil and G.M. Park, 2006. Antiprotozoal activity of deacetylated chitosan oligosaccharide (dp 2-8) on *Trichomonas vaginalis*. *J. Microbiol. Biotechnol.*, 16: 1984-1989.
75. Tavassoli, M., A. Imani, H. Tajik, M. Moradi and S.H. Pourseyed, 2012. Novel *In vitro* efficiency of chitosan biomolecule against *Trichomonas gallinae*. *Iran. J. Parasitol.*, 7: 92-96.
76. Benhamou, N., P.J. Lafontaine and M. Nicole, 1994. Induction of systemic resistance to *Fusarium* crown and root rot in tomato plants by seed treatment with chitosan. *Phytopathology*, 84: 1432-1444.
77. Benhamou, N., J.W. Kloepper and S. Tuzun, 1998. Induction of resistance against *Fusarium* wilt of tomato by combination of chitosan with an endophytic bacterial strain: Ultrastructure and cytochemistry of the host response. *Planta*, 204: 153-168.
78. Bhaskara Reddy, M.V., J. Arul, P. Angers and L. Couture, 1999. Chitosan treatment of wheat seeds induces resistance to *Fusarium graminearum* and improves seed quality. *J. Agric. Food Chem.*, 47: 1208-1216.
79. Benhamou, N. and G. Theriault, 1992. Treatment with chitosan enhances resistance of tomato plants to the crown and root rot pathogen *Fusarium oxysporum* f. sp. *radicis-lycopersici*. *Physiol. Mol. Plant Pathol.*, 41: 33-52.
80. Laflamme, P., N. Benhamou, N. Bussieres and M. Dessureault, 2000. Differential effect of chitosan on root rot fungal pathogens in forest nurseries. *Can. J. Bot.*, 77: 1460-1468.
81. Hadwiger, L.A. and J.M. Beckman, 1980. Chitosan as a component of pea-*Fusarium solani* interactions. *Plant Physiol.*, 66: 205-211.
82. Lizarraga-Paulin, E.G., I. Torres-Pacheco, E. Moreno-Martinez and S.P. Miranda-Castro, 2011. Chitosan application in maize (*Zea mays*) to counteract the effects of abiotic stress at seedling level. *Afr. J. Biotechnol.*, 10: 6439-6446.

83. O'Herlihy, E.A., E.M. Duffy and A. C. Cassells, 2003. The effects of arbuscular mycorrhizal fungi and chitosan sprays on yield and late blight resistance in potato crops from microplants. *Folia Geobotanica*, 38: 201-207.
84. Cho, Y.H., G.J. Choi, B.S. Kim, K.S. Jang, M.Y. Yoon, M.S. Park and J.C. Kim, 2011. Control of late blight of tomato and potato by oilgochitosan. *Res. Plant Dis.*, 17: 129-135.
85. Eikemo, H., A. Stensvand and A.M. Tronsmo, 2003. Induced resistance as a possible means to control diseases of strawberry caused by *Phytophthora* spp. *Plant Dis.*, 87: 345-350.
86. Kim, K.D., S. Nemeč and G. Musson, 1997. Control of *Phytophthora* root and crown rot of bell pepper with composts and soil amendments in the greenhouse. *Applied Soil Ecol.*, 5: 169-179.
87. Falcon, A.B., J.C. Cabrera, D. Costales, M.A. Ramirez, G. Cabrera, V. Toledo and M.A. Martinez-Tellez, 2008. The effect of size and acetylation degree of chitosan derivatives on tobacco plant protection against *Phytophthora parasitica* nicotianae. *World J. Microbiol. Biotechnol.*, 24: 103-112.
88. Liu, H., W. Tian, B. Li, G. Wu and M. Ibrahim *et al.*, 2012. Antifungal effect and mechanism of chitosan against the rice sheath blight pathogen, *Rhizoctonia solani*. *Biotechnol. Lett.*, 34: 2291-2298.
89. Kurzwinska, H., 2007. Potential use of chitosan in the control of lettuce pathogens. Polish Chitin Society, Monograph XII, pp: 173-178.
90. Molloy, C., L.H. Cheah and J.P. Koolaard, 2004. Induced resistance against *Sclerotinia sclerotiorum* in carrots treated with enzymatically hydrolysed chitosan. *Postharvest Biol. Technol.*, 33: 61-65.
91. Yin, H., Y. Li, H.Y. Zhang, W.X. Wang and H. Lu *et al.*, 2013. Chitosan oligosaccharides-triggered innate immunity contributes to oilseed rape resistance against *Sclerotinia sclerotiorum*. *Int. J. Plant. Sci.*, 174: 722-732.
92. El Ghaouth, L.A., J. Arul, N. Benhamou, A. Asselin and R.R. Belanger, 1994. Effect of chitosan on cucumber plants: Suppression of *Pythium aphanidermatum* and induction of defense reactions. *Phytopathology*, 84: 313-320.
93. Anusuya, S. and M. Sathiyabama, 2014. Effect of chitosan on rhizome rot disease of turmeric caused by *Pythium aphanidermatum*. *ISRN Biotechnol.* 10.1155/2014/305349
94. Chien, P.J., F. Sheu and H.R. Lin, 2007. Coating citrus (Murcott tangor) fruit with low molecular weight chitosan increases postharvest quality and shelf life. *Food Chem.*, 100: 1160-1164.
95. Wang, L., H. Wu, G. Qin and X. Meng, 2014. Chitosan disrupts *Penicillium expansum* and controls postharvest blue mold of jujube fruit. *Food Control*, 41: 56-62.
96. Sathiyabama, M. and R. Balasubramanian, 1998. Chitosan induces resistance components in *Arachis hypogaea* against leaf rust caused by *Puccinia arachidis* Speg. *Crop Protec.*, 17: 307-313.
97. Ben-Shalom, N., R. Ardi, R., Pinto, C. Aki and E. Fallik, 2003. Controlling gray mould caused by *Botrytis cinerea* in cucumber plants by means of chitosan. *Crop Protect.*, 22: 285-290.
98. Xu, W.T., K.L. Huang, F. Guo, W. Qu, J.J. Yang, Z.H. Liang and Y.B. Luo, 2007. Postharvest grapefruit seed extract and chitosan treatments of table grapes to control *Botrytis cinerea*. *Postharvest Biol. Technol.*, 46: 86-94.
99. Zhang, D., H. Wang, Y. Hu and Y. Liu, 2015. Chitosan controls postharvest decay on cherry tomato fruit possibly via the mitogen-activated protein kinase signaling pathway. *J. Agric. Food Chem.*, 63: 7399-7404.
100. Bautista-Banos, S., M. Hernandez-Lopez, E. Bosquez-Molina and C.L. Wilson, 2003. Effects of chitosan and plant extracts on growth of *Colletotrichum gloeosporioides*, anthracnose levels and quality of papaya fruit. *Crop Protect.*, 22: 1087-1092.
101. Munoz, Z., A. Moret and S. Garces, 2009. Assessment of chitosan for inhibition of *Colletotrichum* sp. on tomatoes and grapes. *Crop Protect.*, 28: 36-40.
102. Di Piero, R.M. and M.V. Garda, 2008. Chitosan reduces the anthracnose severity and increases the glucanase activity in common bean plants. *Pesquisa Agropecuaria Brasileira*, 43: 1121-1128.
103. Chang, T.H., 2009. Disease control efficacy of chitosan preparations against tomato leaf mold. *Res. Plant Dis.*, 15: 248-253.
104. Cuero, R.G., E. Duffus, G. Osuji and R. Pettit, 1991. Aflatoxin control in preharvest maize: Effects of chitosan and two microbial agents. *J. Agric. Sci.*, 117: 165-169.
105. Dos Santos, N.S.T., A.J.A.A. Aguiar, C.E.V. de Oliveira, C.V. de Sales and S. de Melo e Silva *et al.*, 2012. Efficacy of the application of a coating composed of chitosan and *Origanum vulgare* L. essential oil to control *Rhizopus stolonifer* and *Aspergillus niger* in grapes (*Vitis labrusca* L.). *Food Microbiol.*, 32: 345-353.
106. Yang, L.Y., J.L. Zhang, C.L. Bassett and X.H. Meng, 2012. Difference between chitosan and oligochitosan in growth of *Monilinia fructicola* and control of brown rot in peach fruit. *LWT-Food Sci. Technol.*, 46: 254-259.
107. Manjunatha, G., K.S. Roopa, G.N. Prashanth and H.S. Shetty, 2008. Chitosan enhances disease resistance in pearl millet against downy mildew caused by *Sclerospora graminicola* and defence-related enzyme activation. *Pest Manage. Sci.*, 64: 1250-1257.
108. Maia, A.J., R.V. Botelho, C.M.D.R. Faria and C.D. Leite, 2010. Chitosan action on *Plasmopara viticola* and *Elsinoe ampelina* development *In vitro* and in grapevines cv. Isabel. *Summa Phytopathologica*, 36: 203-209.
109. Algam, S.A.E., G.L. Xie, B. Li, S.H. Yu, T. Su and J. Larsen, 2010. Effects of *Paenibacillus* strains and chitosan on plant growth promotion and control of Ralstonia wilt in tomato. *J. Plant Pathol.*, 92: 593-600.

110. Coqueiro, D.S.O. and M.R. di Piero, 2011. Antibiotic activity against *Xanthomonas gardneri* and protection of tomato plants by chitosan. J. Plant Pathol., 93: 337-344.
111. Ramkissoon, A., J. Francis, V. Bowrin, R. Ramjegathesh, A. Ramsuhag and J. Jayaraman, 2016. Bio efficacy of a chitosan based elicitor on *Alternaria solani* and *Xanthomonas vesicatoria* infections in tomato under tropical conditions. Ann. Applied Biol., 169: 274-283.
112. Li, B., B. Liu, C. Shan, M. Ibrahim and Y. Lou *et al.*, 2013. Antibacterial activity of two chitosan solutions and their effect on rice bacterial leaf blight and leaf streak. Pest Manage. Sci., 69: 312-320.
113. Acar, O., C. Aki and H. Erdugan, 2008. Fungal and bacterial diseases control with Elexa™ plant booster. Fresenius Environ. Bull., 17: 797-802.
114. Pospieszny, H. S. Chirkov and J. Atabekov, 1991. Induction of antiviral resistance in plants by chitosan. Plant Sci., 79: 63-68.
115. Iriti, M., M. Sironi, S. Gomasasca, A.P. Casazza, C. Soave and F. Faoro, 2006. Cell death-mediated antiviral effect of chitosan in tobacco. Plant Physiol. Biochem., 44: 893-900.
116. Nunes da Silva, M., A.R. Cardoso, D. Ferreira, M. Brito, M.E. Pintado and M.W. Vasconcelos, 2014. Chitosan as a biocontrol agent against the pinewood nematode (*Bursaphelenchus xylophilus*). For. Pathol., 44: 420-423.
117. Xing, K., X. Zhu, X. Peng and S. Qin, 2015. Chitosan antimicrobial and eliciting properties for pest control in agriculture: A review. Agron. Sustain. Dev., 35: 569-588.
118. Romanazzi, G., 2010. Chitosan treatment for the control of postharvest decay of table grapes, strawberries and sweet cherries. Fresh Produce, 4: 111-115.
119. Li, P., Z. Cao, Z. Wu, X. Wang and X. Li, 2016. The effect and action mechanisms of oligochitosan on control of stem dry rot of *Zanthoxylum bungeanum*. Int. J. Mol. Sci., Vol. 17. 10.3390/ijms17071044
120. Sudarshan, N.R., D.G. Hoover and D. Knorr, 1992. Antibacterial action of chitosan. Food Biotechnol., 6: 257-272.
121. Helander, I.M., E.L. Nurmiho-Lassila, R. Ahvenainen, J. Rhoades and S. Roller, 2001. Chitosan disrupts the barrier properties of the outer membrane of gram-negative bacteria. Int. J. Food Microbiol., 71: 235-244.
122. Chopra, H. and G. Ruhi, 2016. Eco friendly chitosan: An efficient material for water purification. Pharm. Innov. J., 5: 92-95.
123. Chung, Y.C. and C.Y. Chen, 2008. Antibacterial characteristics and activity of acid-soluble chitosan. Bioresour. Technol., 99: 2806-2814.
124. Hadwiger, L.A., 2013. Multiple effects of chitosan on plant systems: Solid science or hype. Plant Sci., 208: 42-49.
125. Qin, Y., 1993. The chelating properties of chitosan fibers. J. Applied Polym. Sci., 49: 727-731.
126. Rabea, E.I., M.E.T. Badawy, C.V. Stevens, G. Smagghe and W. Steurbaut, 2003. Chitosan as antimicrobial agent: Applications and mode of action. Biomacromolecules, 4: 1457-1465.
127. Bhuvaneshwari, S. and V. Sivasubramanian, 2013. Comparative studies for Chitosan yield and chelating ability of *Aspergillus niger* and *Rhizopus oryzae*. Indian J. Biotechnol., 12: 429-431.
128. Freeman, B.C. and G.A. Beattie, 2008. An overview of plant defenses against pathogens and herbivores. Plant Health Instructor, 10.1094/PHI-I-2008-0226-01
129. Lin, W., X. Hu, W. Zhang, W.J. Rogers and W. Cai, 2005. Hydrogen peroxide mediates defence responses induced by Chitosans of different molecular weights in rice. J. Plant Physiol., 162: 937-944.
130. Eilenberg, H., S. Pnini-Cohen, Y. Rahamim, E. Sionov, E. Segal, S. Carmeli and A. Zilberstein, 2009. Induced production of antifungal naphthoquinones in the pitchers of the carnivorous plant *Nepenthes khasiana*. J. Exp. Bot., 61: 911-922.
131. Burkhanova, G.F., L.G. Yarullina and I.V. Maksimov, 2007. The control of wheat defense responses during infection with *Bipolaris sorokiniana* by chitooligosaccharides. Russian J. Plant Physiol., 54: 104-110.
132. Hoat, T.X., H. Nakayashiki, Q. Yang, Y. Tosa and S. Mayama, 2013. Molecular cloning of the apoptosis-related calcium-binding protein AsALG-2 in *Avena sativa*. Mol. Plant Pathol., 14: 222-229.
133. Fan, G., X. Li, X. Wang, Q. Zhai and Y. Zhan, 2010. Chitosan activates defense responses and triterpenoid production in cell suspension cultures of *Betula platyphylla* suk. Afr. J. Biotechnol., 9: 2816-2820.
134. Katiyar, D., A. Hemantaranjan and B. Singh, 2015. Chitosan as a promising natural compound to enhance potential physiological responses in plant: A review. Indian J. Plant Physiol. 10.1007/s40502-015-0139-6
135. Klessig, D.F., J. Durner, J. Shah and Y. Yang, 1998. Salicylic Acid-Mediated Signal Transduction in Plant Disease Resistance. In: Phytochemical Signals and Plant-Microbe Interactions, Romeo, T.J., R.K. Downum and R. Verpoorte (Eds.). Springer, New York, pp: 119-137.
136. Petutschnig, E.K., A.M. Jones, L. Serazetdinova, U. Lipka and V. Lipka, 2010. The Lysin Motif Receptor-Like Kinase (LysM-RLK) CERK1 is a major chitin-binding protein in *Arabidopsis thaliana* and subject to chitin-induced phosphorylation. J. Biol. Chem., 285: 28902-28911.
137. Lienart, Y., C. Gautier and A. Domard, 1991. Isolation from *Rubus* cell-suspension cultures of a lectin specific for glucosamine oligomers. Planta, 184: 8-13.
138. Malerba, M. and R. Cerana, 2016. Chitosan effects on plant systems-A review. Int. J. Mol. Sci., Vol. 17, No. 7. 10.3390/ijms17070996

139. EPA., 2003. Chitosan; Poly-D-glucosamine (128930) fact sheet. Environmental Protection Agency (EPA), Washington, DC., USA., June 1, 2003.
140. Cretoiu, M.S., G.W. Korthals, J.H. Visser and J.D. van Elsas, 2013. Chitin amendment increases soil suppressiveness toward plant pathogens and modulates the actinobacterial and oxalobacteraceal communities in an experimental agricultural field. *Applied Environ. Microbiol.*, 79: 5291-5301.
141. Mulawarman, J. Hallmann, D. Bell, B. Kopp-Holtwiesche and R.A. Sikora, 2001. Effects of natural products on soil organisms and plant health enhancement. *Meded. Rijksuniv. Gent. Fak. Landbouwk. Toegep. Biol. Wet.*, 66: 609-617.
142. Pastucha, A., 2005. The effect of chitosan on the formation of microorganism communities in the rhizosphere soil of soybean. *Acta Scientiarum Polonorum-Hortorum Cultus*, 4: 69-77.
143. Postma, J., L.H. Stevens, G.L. Wieggers, E. Davelaar and E.H. Nijhuis, 2009. Biological control of *Pythium aphanidermatum* in cucumber with a combined application of *Lysobacter enzymogenes* strain 3.1T8 and chitosan. *Biol. Control*, 48: 301-309.
144. Manjula, K. and A.R. Podile, 2001. Chitin-supplemented formulations improve biocontrol and plant growth promoting efficiency of *Bacillus subtilis* AF 1. *Can. J. Microbiol.* 47: 618-625.
145. Lowe, A., S.M. Rafferty-McArdle and A.C. Cassells, 2012. Effects of AMF- and PGPR-root inoculation and a foliar chitosan spray in single and combined treatments on powdery mildew disease in strawberry. *Agric. Food Sci.*, 21: 28-38.
146. Mishra, S., K.S. Jagadeesh, P.U. Krishnaraj and S. Prem, 2014. Biocontrol of tomato leaf curl virus (ToLCV) in tomato with chitosan supplemented formulations of *Pseudomonas* sp. under field conditions. *Aust. J. Crop Sci.*, 8: 347-355.
147. Staehelin, C., M. Schultze, K. Tokuyasu, V. Poinso, J.C. Prome, E. Kondorosi and A. Kondorosi, 2000. N-deacetylation of *Sinorhizobium meliloti* Nod factors increases their stability in the *Medicago sativa* rhizosphere and decreases their biological activity. *Mol. Plant-Microbe Interact.*, 13: 72-79.
148. Genre, A., M. Chabaud, C. Balzergue, V. Puech-Pages and M. Novero *et al.*, 2013. Short-chain chitin oligomers from arbuscular mycorrhizal fungi trigger nuclear Ca²⁺ spiking in *Medicago truncatula* roots and their production is enhanced by strigolactone. *New Phytol.*, 198: 190-202.
149. Walker, T.S., H.P. Bais, E. Grotewold and J.M. Vivanco, 2003. Root exudation and rhizosphere biology. *Plant Physiol.*, 132: 44-51.
150. Bais, H.P., T.L. Weir, L.G. Perry, S. Gilroy and J.M. Vivanco, 2006. The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu. Rev. Plant Biol.*, 57: 233-266.
151. Zeng, D. and D. Luo, 2012. Physiological effects of chitosan coating on wheat growth and activities of protective enzyme with drought tolerance. *Open J. Soil Sci.*, 2: 282-288.
152. Herde, O., H. Pena-Cortes, L. Willmitzer and J. Fisahn, 1997. Stomatal responses to jasmonic acid, linolenic acid and abscisic acid in wild-type and ABA-deficient tomato plants. *Plant. Cell. Environ.*, 20: 136-141.
153. Bittelli, M., M. Flury, G.S. Campbell and E.J. Nichols, 2001. Reduction of transpiration through foliar application of chitosan. *Agric. Forest Meteorol.*, 107: 167-175.
154. Pongprayoon, W., S. Roytrakul, R. Pichayangkura and S. Chadchawan, 2013. The role of hydrogen peroxide in chitosan-induced resistance to osmotic stress in rice (*Oryza sativa* L.). *Plant Growth Regul.*, 70: 159-173.
155. Je, J.Y. and S.K. Kim, 2006. Reactive oxygen species scavenging activity of aminoderivatized chitosan with different degree of deacetylation. *Bioorg. Med. Chem.*, 14: 5989-5994.
156. Rajalakshmi, A., N. Krithiga and A. Jayachitra, 2013. Antioxidant activity of the chitosan extracted from shrimp exoskeleton. *Middle-East J. Scient. Res.*, 16: 1446-1451.
157. Kamari, A., I.D. Pulford and J.S.J. Hargreaves, 2011. Binding of heavy metal contaminants onto chitosans-An evaluation for remediation of metal contaminated soil and water. *J. Environ. Manage.*, 92: 2675-2682.
158. Sanchez-Duarte, R.G., D.I. Sanchez-Machado, J. Lopez-Cervantes and M.A. Correa-Murrieta, 2012. Adsorption of allura red dye by cross-linked chitosan from shrimp waste. *Water Sci. Technol.*, 65: 618-623.
159. Yong, S.K., M. Shrivastava, P. Srivastava, A. Kunhikrishnan and N. Bolan, 2015. Environmental applications of chitosan and its derivatives. *Rev. Environ. Contam. Toxicol.*, 233: 1-43.